



# GRANULAR SYSTEMIC FUNGICIDE

Contains Thiophanate-methyl

- Prevents or arrests the turf grass diseases, brown patch, dollar spot, and copper spot.
- Application flexibility – may be applied to dry or moist foliage using a drop-type or rotary spreader.
- Fast acting, sustained disease control through root and foliage uptake, resulting in less susceptibility of product to weather and maintenance conditions.
- Choice of application rates, depending upon disease, severity of infection and kind of protection desired.

## GENERAL INFORMATION

Bulk Density:	27.0 lb/cu. ft.
Net Weight:	13.49 kg (29.81 lb)
Coverage:	2,044 m <sup>2</sup> (22,000 sq. ft.)

## PRODUCT SPECIFICATIONS

**Active ingredient:** Thiophanate-methyl 2.3%

A dry, ready to use, free flowing product with active ingredient impregnated on and into a granular carrier.

# GRANULAR SYSTEMIC FUNGICIDE

Contains Thiophanate-methyl

REGISTRATION NUMBER 16660 PEST CONTROL PRODUCTS ACT

**FOR PROFESSIONAL USE ONLY.**  
**KEEP OUT OF REACH OF CHILDREN.**  
**READ THE LABEL BEFORE USING.**

## DIRECTIONS FOR USE:

Prevents or arrests the turfgrass diseases brown patch, dollar spot, and copper spot.

**NORMAL RATE:** 6.6 kg / 1,000 m<sup>2</sup>

**DOUBLE RATE:** 13.2 kg / 1,000 m<sup>2</sup>

For prevention of dollar spot, copper spot, and brown patch, apply to moist or dry foliage at the **NORMAL RATE** when disease activity is noted or suspected and repeat at 2 to 3 week intervals. Brown patch control may require more frequent treatments under conditions especially favourable to disease activity.

To arrest these diseases, apply at the **DOUBLE RATE** and follow with the **NORMAL RATE** preventative program.

## PRECAUTIONS:

**KEEP OUT OF REACH OF CHILDREN.** May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Do not contaminate feed or foodstuff. Keep children and pets off treated area until material has been washed off the foliage and area is completely dry. Do not feed clippings from treated turf to livestock and poultry. Do not contaminate domestic or irrigation water supplies, lakes, streams and ponds.

## STORAGE:

Store in cool, dry, ventilated place. Keep away from heat, fire and sparks. Wash thoroughly before eating, drinking or smoking. Wash contaminated clothing with soap and hot water before reuse.

## DISPOSAL:

1. Thoroughly empty the contents of the container into the application device.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## FIRST AID:

**IF ON SKIN,** remove contaminated clothing and wash skin thoroughly with soap and water. **IF IN EYES,** flush with water for 5-10 minutes and obtain medical attention **IMMEDIATELY.** **IF SWALLOWED,** give patient 1 to 2 glasses of water and cause vomiting by inserting a finger or blunt object down the throat. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Obtain medical attention or contact a poison control centre **IMMEDIATELY.** Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION:

Treat symptomatically.

## NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

## NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

## SUGGESTED SPREADER SETTINGS

To provide proper distribution, calibrate spreader before application

13.49 kg treats 2,044 m<sup>2</sup> at the **NORMAL RATE** (Use 6.6 kg/1,000 m<sup>2</sup>)

13.49 kg treats 1,022 m<sup>2</sup> at the **DOUBLE RATE** (Use 13.2 kg/1,000 m<sup>2</sup>)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			NORMAL RATE	DOUBLE RATE
ProTurf® Drop Type	5 km/h	Overlap wheels	4 3/4	6
ProTurf® Rotaries (R-7, R-7X) R-8 (cone 8)	5 km/h	2.4 m	C	F
	5 km/h	2.7 m	J	L
Accupro 2000 (cone 8)	5 km/h	2.7 m	1 1/2	K
SR 2000 (cone 8)	5 km/h	2.7 m	1 1/2	K
Lely Models WTR, WFR, HR and 1250. PTO at 450 rpm	7 km/h	4.9 m	3 1/2 III	4 1/2 III



Always refer to the actual product label for complete directions and precautions.

ProTurf® is a registered trademark, used under licence from The Scotts Company. Nu-Gro IP Inc., 10 Craig Street, Brantford, Ontario N3R 7J1