

## SAFER'S INSECTICIDAL SOAP CONCENTRATE

For control of aphids, mealybugs, spider mites, whitefly, soft brown scale, psyllids, rose or pear slugs (sawfly larvae), and earwigs on houseplants, roses, flowers, vegetables, fruits, ornamentals, shrubs, trees or greenhouse plantings; in or around the house or home.

Kills insects on contact. Can be used indoors or outdoors.

GUARANTEE: Soap (Potassium salts of Fatty Acids) .....50.5%

REG. NO. 22208 P.C.P. ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

NET CONTENTS     mL, L

### DOMESTIC

Woodstream Canada Corporation, 4 Lowry Drive, Brampton, ON L7A 1C4  
(905) 840-2640

**DIRECTIONS:** Mix fresh solution. Made of selected soaps that have contact insecticidal properties. Insects must be sprayed directly to achieve control. Mix one part concentrate to 50 parts water, or 10 mL per 500 mL water. **Spraying:** Spray stems and both leaf surfaces thoroughly. **Dipping:** (for small potted plants): Pad the soil with paper or plastic to prevent spillage, turn the plant upside down and dip stems and leaves in dilute solution for three seconds. Product may be used on fruit and vegetables up to and including the day of harvest. Wash fruits and vegetables before eating.

### INSECTS CONTROLLED:

**APHIDS:** Aphids can fly outdoors from early spring throughout the summer. Aphids can occur year round on plants indoors. Spray when sighted on buds, shoots, stems and the undersides of leaves. Aphids must be contacted to be controlled. Repeat applications (spray) weekly for 2-3 weeks and thereafter as required.

**MEALYBUGS:** Found on all plant parts. Spray when sighted on buds, shoots, stems and the undersides of leaves. Mealybugs must be contacted to be controlled. Apply (spray) weekly for 2-3 weeks and thereafter as required.

**WHITEFLY:** These tiny moth-like insects can be found throughout the year on the undersides of indoor plants and throughout the growing season on outdoor plantings on buds, shoots, stems and the undersides of leaves. Apply (spray) weekly for 2-3 weeks and thereafter as required to control the scale-like larvae hatching from eggs and adult flies.

**SPIDER MITES:** Mites are tiny spider-like creatures with 8 legs which feed on plant juices causing yellowing and browning of leaves. Spider mites are often first observed

by the webs they spin on the leaves and buds of plants. Apply (spray) weekly for 2-3 weeks and thereafter as required. Foliage should also be misted with water daily until mite control is achieved.

**EARWIGS:** Earwigs are dark brown crawling insects approximately 2.5 cm (1 inch) in length. They can easily be identified by the pincers that extend from the rear of the body. Earwigs hide in plants, garden mulch, and crevices during the day and feed at night. They feed on a wide variety of plants leaving holes in the foliage. For control, spray at dusk or in the evening in all areas where earwig damage is evident. Spray all plants thoroughly including the soil. If spraying at night, a light may be used to identify infestations. Repeat applications as required.

**SOFT BROWN SCALE:** Scale crawlers are killed on contact. Staggered hatching of crawlers requires 2-3 spray applications to achieve control. Spray or apply at 7-10 day intervals as required.

**ROSE AND PEAR SLUGS (SAWFLY LARVAE):** Spray both leaf surfaces to achieve control. Insects must be sprayed directly to achieve control. Watch for a second generation to appear and repeat applications as required.

**PRECAUTIONS:** Concentrated soap may cause eye irritation. Avoid eye contact. May cause skin irritation. Do not use on sweetpeas, nasturtiums, delicate ferns or impatiens. If in doubt about a sensitive species, perform a small-scale test spray before treating entire plant. Do not apply to plants in direct sunlight. To maximize blossom longevity, avoid spraying blooms during the heat of the day.

**DISPOSAL:** Do not reuse empty container. Wrap empty container and discard in household garbage.

**FIRST AID:** If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. If ingested, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically

The active ingredients in this product are specially selected soaps. No other pesticides are added.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.