

PLANT PRODUCTS SAFETRAY™ P

Wood Preservative for the Mushroom Industry

For the control of adhesion and penetration of mushroom mycelium in timber trays and benches allowing for an easy tip out of the adherent compost.

**COMMERCIAL
KEEP OUT OF REACH OF CHILDREN**

CAUTION POISON

EYE IRRITANT

READ THE LABEL BEFORE USING

GUARANTEE:

Propiconazole* 242.5 g/litre

NET CONTENTS: 10 L

REGISTRATION NO.: 24813.01
PEST CONTROL PRODUCTS ACT

™ Janssen Pharmaceutica, Beerse, Belgium.

* Active ingredient and formulation developed by Janssen Pharmaceutica, Beerse, Belgium

PLANT PRODUCTS CO. LTD.
314 Orenda Road
Brampton, Ontario
L6T 1G1

DIRECTIONS FOR USE: For the control of adhesion and penetration of mushroom mycelium in timber trays and benches add 6 ml. of Safetray™ P to 1 litre of water to achieve a treating concentration of 1,500 ppm active ingredient.

MIXING: Half-fill the spray tank with water. Add the appropriate amount of SAFETRAY™ P. Triple rinse empty container and add the rinsate to the tank. Stir gently and add water up to the required volume. A few drops of an anti-foam agent may be added to avoid excessive foaming.

APPLICATION: Apply the spray solution with a large droplet sprayer on the internal surface of the trays or to the inside surface of the side, end and bottom boards of the benches. To insure complete wetting of the wood surface use approximately 95 litres of spray solution per 465 m² (5,000 sq. ft.) of wood surface. Make sure all corners are well treated. Let the trays or bench boards dry before adding compost. The trays should be allowed to dry in a sheltered, well-ventilated place.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through the skin. Avoid contact with skin or eyes. Avoid breathing spray mist. Wear coveralls or long pants and long-sleeved shirt, respiratory protection and gloves when mixing and loading. On tray farms, application should be

through mechanical systems. For manual application, wear impermeable suit, gloves and respiratory protection during application. Ventilate prior to reentry. Do not reenter treated area until residues have dried. If treated trays are handled prior to drying, wear gloves. Wash thoroughly with soap and water after handling and before eating and smoking. Avoid contamination of food and feed and domestic or irrigation water supplies, lakes, streams and ponds. Do not reuse container, destroy when empty.

STORAGE: Store in original container in a cool, dry place. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID:

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote known. Treat symptomatically. Avoid giving alcohol or alcohol related products.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Product Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.