

GROUP	40	FUNGICIDE
-------	----	-----------

MICORA™ Fungicide

FUNGICIDE

SUSPENSION

COMMERCIAL

For control or suppression of downy mildew or listed Phytophthora diseases in greenhouse and outdoor ornamental crops, greenhouse tomatoes, bell peppers and lettuce.

GUARANTEE:

Mandipropamid 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO: **30759**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote if this product is ingested. Treat symptomatically. There is no known medical condition likely to be aggravated by the exposure.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Applicators and other handlers must wear long-sleeve shirt and long pants when mixing, loading and applying and during clean-up and repair activities. Wear chemical resistant gloves during mixing, loading, clean-up and repair activities. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind

speed, wind direction, temperature inversion, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

ENVIRONMENTAL PRECAUTIONS

Runoff

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

Store in a cool, dry place. Do not store food, beverages or tobacco products in storage area.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (1-800-FASTMED)***

GENERAL INFORMATION

MICORA™ Fungicide is a fungicide for control or suppression of diseases in greenhouse tomatoes and greenhouse lettuce.

DIRECTIONS FOR USE

Rotational Crop Restrictions

Do not plant any crop which is not registered for use with MICORA Fungicide for a period of 30 days after the last application.

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), estuarine/marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters .

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use control aquatic pests.

APPLICATION PROCEDURES

Ground Application:

Spray nozzles should be selected which will provide accurate and uniform spray deposition. Use spray nozzles, which provide medium-sized droplets and reduce drift. To help ensure accuracy, calibrate sprayer before each use. Do not apply MICORA Fungicide through any type of ultra low volume (ULV) spray system. For information on spray equipment and calibration, consult sprayer manufacturers and/or the local agricultural representative specialists.

Apply MICORA Fungicide using sufficient water volume to provide thorough and uniform coverage. Use a minimum of 100 L/ha unless otherwise indicated on this label. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. Avoid making applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. Avoid spray overlap.

Do not apply this product using fogging equipment (handheld or automated), or using handheld mist blowers/airblast equipment.

Precautions:

- 1) Select nozzles that can produce the desired droplet sizes at the normal rated pressure range.
- 2) The sprayer should travel at a uniform speed across the crop.

Sprayer application: **DO NOT** apply during periods of dead calm. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. Boom height must be 60 cm or less above the crop or ground.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-87-SYNGENTA (1-877-964-3682) or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in the greenhouse water volume guidelines.

Greenhouse Water Volume Guidelines

Crop	Size	Height (m)	Volume per unit area (L/ha)
Tomato	Small	0.6	285
	Medium	1.2	627
	Large	2.7	1200-1400
Ornamentals	Average ornamental crop		1000

MICORA Rate Conversion Chart

mL Product/ha	g ai/ha	Treated ha/1 L Product
400	100	2.5
500	125	2.0
600	150	1.7

MIXING PROCEDURES

1. Add 1/2 of the required amount of water to the mix tank.
2. With the agitator running, add the desired amount of MICORA Fungicide to the tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after MICORA Fungicide has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

DIRECTIONS FOR USE

Integrated Pest (Disease) Management: MICORA Fungicide should be integrated into an overall disease management strategy that includes varieties with disease tolerance, proper timing of irrigation and removal of plant debris in which inoculum overwinters.

Resistance Management: Follow the resistance management guidelines in the Resistance Management Recommendations Section. Do not apply more than one application of MICORA Fungicide or any other Group 40 fungicide before alternation with at least one application of a fungicide that is not a group 40.

GREENHOUSE VEGETABLE CROPS	
CROP	GREENHOUSE TOMATOES (For use in greenhouse only - not for transplant to the field)
PEST	For control of Late blight (<i>Phytophthora infestans</i>)
RATE	400-600 mL product/ha (100-150 g ai/ha)
APPLICATION INTERVAL (days)	7-10
PHI (days)	1
APPLICATION DIRECTIONS	MICORA Fungicide applications should begin prior to disease development and continue throughout the season on a 7-10 day schedule of fungicides, following the resistance management guidelines. The use of a non-ionic adjuvant (0.25% v/v) is recommended. Refer to the Greenhouse Water Volumes chart. DO NOT make more than four applications per crop cycle of MICORA Fungicide.

CROPS	GREENHOUSE LETTUCE (For use in greenhouse only - not for transplant to the field)
PESTS	For control of Downy mildew (<i>Bremia lactucae</i>), Blue Mold (<i>Peronospora effusa</i>)
RATE	400-600 mL product/ha (100-150 g ai/ha)
APPLICATION INTERVAL (days)	7-10
PHI (days)	7
APPLICATION DIRECTIONS	MICORA Fungicide applications should begin prior to disease development and continue throughout the season on a 7-10 day schedule of fungicides, following the resistance management guidelines. The use of a non-ionic adjuvant (0.25% v/v) is recommended. DO NOT make more than four applications per crop cycle of MICORA Fungicide.

READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS

NOTE TO USER:

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Syngenta Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Syngenta Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerances arising, and agree to hold Syngenta Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

GREENHOUSE AND OUTDOOR ORNAMENTALS	
CROP	GREENHOUSE AND OUTDOOR ORNAMENTALS
PEST	Control of Downy Mildew caused by <i>Peronospora spp.</i> Suppression of Phytophthora foliar blight caused by <i>Phytophthora nicotianae</i> and <i>Phytophthora ramorum</i>
RATE	30 – 60 mL/100L
APPLICATION INTERVAL (days)	7-14 days

APPLICATION DIRECTIONS	<p>Begin foliar applications prior to disease development and continue on a 7- to 14- day interval when conditions are favorable for disease development as a preventative spray or at first sign of disease symptoms.</p> <p>Make no more than 2 consecutive applications before switching to another effective non-Group 40 fungicide.</p> <p>Use the shorter interval and/or higher rates under high disease pressure or when conditions are conducive to disease development.</p> <p>The use of a non-ionic adjuvant (0.25% v/v) is recommended.</p> <p>DO NOT make more than four applications per crop cycle of MICORA Fungicide.</p> <p>Plant safety has been found to be acceptable for many ornamental plants, however, not all possible plant species and varieties have been tested under all conditions. It is recommended to test MICORA Fungicide on a small portion of the crop to ensure that a phytotoxic response will not occur.</p>
CROPS	GREENHOUSE BELL PEPPER (For use in greenhouse only - not for transplant to the field)
PESTS	For suppression of root and crown rot caused by <i>Phytophthora capsici</i>
APPLICATION METHOD	Apply as a drench to the base of the plant in sufficient water to ensure the root area is covered.
PESTS	For suppression of Phytophthora blight on the stems, leaves and fruit caused by <i>Phytophthora capsici</i>
APPLICATION METHOD	Apply as a foliar spray in sufficient water to ensure good coverage of foliar tissue. Refer to Greenhouse Water Volumes Guidelines.
RATE	600 mL product/ha (150 g ai/ha)
APPLICATION INTERVAL	7-10 days
PREHARVEST INTERVAL (PHI)	1 day
APPLICATION DIRECTIONS	<p>MICORA Fungicide applications should begin prior to disease development and continue throughout the season on a 7-10 day schedule of fungicides, following the resistance management guidelines.</p> <p>DO NOT make more than 1 drench application and 3 foliar applications of MICORA Fungicide per crop cycle.</p>

RESISTANCE MANAGEMENT RECOMMENDATIONS

For Resistance management, please note that MICORA Fungicide contains a Group 40 fungicide. Any fungal population may contain individuals naturally resistant to MICORA Fungicide and other Group 40 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same crop. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of MICORA Fungicide or other Group 40 fungicides with different Groups that control the same pathogens.

Do not exceed the total number of applications of MICORA Fungicide per season per crop.

Do not apply sequential treatments of MICORA Fungicide, or other fungicides in the same Fungicide Group, in a season. Do not apply at rates lower than recommended on the label.

Use tank mixtures with fungicides from a different Group when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications. Monitor treated fungal populations for sign of resistance development.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information and to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

MICORA™ is a trademark of a Syngenta Group Company.

GROUP	40	FUNGICIDE
-------	----	-----------

MICORA™ Fungicide

FUNGICIDE

SUSPENSION

COMMERCIAL

For control or suppression of downy mildew or listed Phytophthora diseases in greenhouse and outdoor ornamental crops, greenhouse tomatoes, peppers and lettuce.

GUARANTEE:

Mandipropamid 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO: 30759
PEST CONTROL PRODUCTS ACT

NET CONTENT: 50 mL to Bulk

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

Label

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote if this product is ingested. Treat symptomatically. There is no known medical condition likely to be aggravated by the exposure.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Applicators and other handlers must wear long-sleeve shirt and long pants when mixing, loading and applying and during clean-up and repair activities. Wear chemical resistant gloves during mixing, loading, clean-up and repair activities. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

ENVIRONMENTAL PRECAUTIONS

Runoff

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

Store in a cool, dry place. Do not store food, beverages or tobacco products in storage area.

DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

3. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
4. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (1-800-FASTMED)***