

GROUP	21	INSECTICIDE
-------	----	-------------

NEXTER® WP Miticide/Insecticide

For control of mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards (Ontario only), mites as a post-harvest application on raspberries, and pear psylla in pears.

COMMERCIAL

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 75%

REGISTRATION NO.: 25135
PEST CONTROL PRODUCTS ACT



WARNING

POISON

Warning, contains the allergen sulfite

**DANGER: FATAL IF INHALED. DO NOT BREATHE DUST, VAPOUR OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES OR CLOTHING.**

**HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN.
READ THE LABEL AND BOOKLET BEFORE USING**

DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY

NET CONTENTS:



Canyon Group, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

GENERAL INFORMATION

This box contains **NEXTER WP Miticide/Insecticide**, a 75% wettable powder, in water-soluble bags (PVA) to control mite pests, including European red mite, apple rust mite, McDaniel spider mite, pear rust mite and twospotted spider mite, when used at recommended rates.

NEXTER WP Miticide/Insecticide works primarily through contact action, so good spray coverage is necessary.

NEXTER WP Miticide/Insecticide provides excellent knockdown and residual control.

DIRECTIONS FOR USE

Please refer to and retain this booklet for complete directions. Do not handle PVA bags excessively or expose them to moisture as this may cause breakage. Bags should not become wet prior to adding to the spray tank. Do not handle the bags with wet hands or wet gloves. Always reseal the overwrap foil envelope to protect the remaining unused bags.

Do not remove water-soluble bags from overwrap except to add them directly to the spray tank.

Do not place the PVA bags directly in diesel oils, dormant or summer-spray-type oils. PVA bags are water soluble, not oil soluble.

Grapes

Apply to grapes when immature motile stages are present for best control.

Pears

Apply when pear psylla numbers reach significant levels. Second application to pears for mites and pear psylla can be made 30 days after the first application.

Raspberries

Apply to raspberry plants after harvest when mites appear..

The use of NEXTER miticide/insecticide may be compatible with the goal of preservation of predatory mites. This product is easiest on predatory mites when applied late spring to early summer before numbers of beneficial mites have peaked. NEXTER miticide/insecticide is harmful to certain beneficials (i.e. beneficial insects).

APPLICATION RATES

Table 1. NEXTER WP Miticide/Insecticide Application Rate Table

Mites Controlled	Rate per hectare	Use Directions
European Red Mite (<i>Panonychus ulmi</i>)	300 grams (2.5 bags)	Apply NEXTER WP in a spray volume of 550-2000 L/ha. Apply in a sufficient water volume to ensure thorough coverage. Nexter WP can only be used on fruit trees for which thorough coverage can be achieved with a maximum spray volume of 2000 L/ha.
Apple Rust Mite (<i>Aculus schlechtendali</i>)		
Pear Rust Mite (<i>Epirimerus pyri</i>)		
Twospotted Spider Mite (<i>Tetranychus urticae</i>)	300* or 600 grams** (2.5 or 5 bags)	DO NOT use NEXTER WP in successive applications. Use NEXTER WP in rotation with other miticides/insecticides with different modes of action.
McDaniel Spider Mite (<i>Tetranychus mcdanieli</i>)		
Pear Psylla (<i>Cacopsylla pyricola</i>)	600 - 720 grams (5 - 6 bags)	

* Good control will be achieved at or below the mite action threshold.

** For faster knockdown and control above mite action thresholds late season

NOTE: NEXTER WP Miticide/Insecticide controls nymph and larva stage of twospotted spider and McDaniel mites and all motile stages of European red, apple rust mite and pear rust mite. This product is not an ovicide.

For foliar application using ground application equipment only.

Crop Recommendation Table

Crop	Application to harvest interval (days)	Maximum number of applications / Year	Maximum product used (kg/ha)/ year on the same orchard	Minimum interval between Nexter WP Applications
Apples	25	2	1.2	30 days
Cherries (sweet and tart)	7	1	0.6	Not applicable.
Grapes	25	1	0.6	Not applicable.
Peaches/Nectarines (Ontario only)	14	1	0.6	Not applicable.
Pears	25	2	1.44	30 days
Raspberries: Post-harvest application only	N/A	2	1.2	30 days
Strawberries	10	2	1.2	15 days

*For optimal control, apply product in sufficient water for good leaf coverage (3000 L/ha).

TIMING

NEXTER WP Miticide/Insecticide should be applied when populations are building and before serious crop injury occurs. Consult provincial guidelines and local extension experts for treatment threshold and advice. Mites cause the most economic damage one week to one month following bloom. **NEXTER WP Miticide/Insecticide** provides excellent knockdown and residual control. A good performance evaluation can be made 7-10 days after treatment.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **NEXTER WP Miticide/Insecticide** contains a Group 21 insecticide/acaricide. Any insect/mite population may contain individuals naturally resistant to **NEXTER WP Miticide/Insecticide** and other Group 21 insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of **NEXTER WP Miticide/Insecticide** or other Group 21 insecticides/acaricides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides/acaricides from a different group when such use is permitted.
- Insecticide/acaricide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

- For further information or to report suspected resistance, contact Gowan Company at 1-800-883-1844.

MIXING

1. Fill the spray tank one-half full of water and start agitation.
2. Add the required number of water-soluble bags to the tank.
3. Continue agitation while filling tank to the desired spray volume.
4. Allow the bags to completely dissolve before spraying.
5. Dilute **NEXTER WP Miticide/Insecticide** in enough water to provide thorough spray coverage. If foaming occurs, an anti-foaming agent may be used.

NOTE: Do not attempt to use the PVA bags directly in diesel, oils or summer-type oils as in ULV or LV use. PVA bags are water-soluble, not oil-soluble.

TANK MIXING

Physical incompatibility, reduced pest control or crop injury may result from mixing **NEXTER WP Miticide/Insecticide** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. Gowan Company does not offer any recommendations for tank mixing **NEXTER WP Miticide/Insecticide** with any other products.

RESTRICTIONS AND LIMITATIONS

Do not use products that contain boron because boron will prevent the water-soluble bags from dissolving. If boron-containing products have been used in previous applications, thoroughly wash the spray tank according to the recommended tank cleaning methods.

Do not apply this product through any type of irrigation system.

DO NOT APPLY THIS PRODUCT BY AIR.

Do not apply directly to aquatic systems. Aquatic systems include all permanent lentic (standing) and lotic (flowing) water bodies. This product is toxic to aquatic invertebrates, fish and aquatic plants. To mitigate impact on these organisms, a minimum buffer zone width of 49 m from aquatic systems should be observed with applications with air-blast sprayers.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 8 km/h at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Air-induction nozzles must be used for the ground application of this product. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Buffer zones:

Spot treatments using hand-held equipment DO NOT require a buffer zone. The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop		Buffer Zones (meters) Required for Protection of:				Terrestrial habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths		
			Less than 1m	Greater than 1 M	Less than 1m	Greater than 1 M	
Groundboom	Strawberry, raspberry		20	2	1	1	1
	Outdoor ornamental		10	1	1	1	0
Airblast	Pear, apple, raspberry	Late growth stage	65	40	30	20	1
	Cherry, peach, nectarine, grapes	Late growth stage	60	35	30	20	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for airblast application of this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site. Buffer zones for groundboom sprayer application CANNOT be modified using the Buffer Zone Calculator.

To reduce run-off from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section.

For apples, raspberries, cherries, nectarines, peaches, pears and outdoor ornamentals: TOXIC to bees. DO NOT apply during the crop blooming period.

For grapes and strawberries: TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

DO NOT apply by hand-held mistblower or hand-held fogger.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

1. May be harmful if swallowed. Do not take internally.
2. May be fatal if inhaled. Do not inhale dust, vapour or spray mist. May irritate eyes. Avoid contact with eyes, skin and clothing.
3. Wash thoroughly after handling and before eating, drinking or smoking.
4. Wear long-sleeved shirt, long pants, rubber boots, socks, goggles, gloves (rubber, PVC, neoprene or nitrile), hat and a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides during mixing, loading and application. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI). Employers should make every effort to schedule pesticide applications and worker tasks in order to avoid early entry of workers into treated areas. Under exceptional circumstances, certified pesticide applicators may enter treated areas for short-term tasks not involving hand labour if at least 4 hours have passed since application and a long sleeved shirt, long pants, rubber boots, socks, goggles, gloves (rubber, PVC, neoprene or nitrile), hat, and a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 24 hour period or until the REI ends.
5. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
6. Clean spray equipment thoroughly after use.
7. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
8. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply or allow **NEXTER WP Miticide/Insecticide** to drift to crops or weeds on which bees are actively foraging. Do not allow spray to drift towards beehives.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca .

This product is only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the pesticide application is to occur.

DO NOT use in residential areas. Residential areas are defined as sites where bystanders including children may be potentially exposed during or after spraying. This includes around homes, school, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed.

Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Hazardous to humans and domestic animals. Keep out of reach of children and pets.

DO NOT allow entry or allow workers to enter treated areas during the restricted entry intervals REI(s) specified in the following table.

Crop	Postapplication Activity	Restricted Entry Interval
Grapes	Girdling, turning	54 Days
	Tying (full foliage), training (full foliage), hand harvest, leaf pulling by hand	30 days
	All other activities (e.g. scouting, weeding, irrigation, bird control, propagating, trellis repair, pruning, transplanting)	12 hours
All other crops	All activities	12 hours

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a major spill, leak or fire call Chemtrec® at 1-800-424-9300. For medical emergencies call 1-888-478-0798

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying - Best Management Practices" on the Health Canada website (www.healthcanada.gc.ca/pollinators). Follow crop specific directions for application timing.

For applications on crops that are highly attractive to pollinators (apples, raspberries, cherries, nectarines, peaches, pears, and outdoor ornamentals), and crops where managed bees are present: DO NOT apply during the crop blooming period.

For applications to grapes and strawberries: Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

TOXIC to beneficial arthropods. Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.
TOXIC to birds.

TOXIC to small wild mammals.

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

STORAGE

1. Store in original tightly closed container. Do not expose water-soluble bags to moisture.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. To prevent contamination store this product away from food or feed

DISPOSAL

1. Follow provincial instructions for any required cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.

4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2019-1328
2019-09-19

Nexter – base panel for booklet – permanently attached to foil bag

GROUP	21	INSECTICIDE
-------	----	-------------

NEXTER® WP Miticide/Insecticide

For control of mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards (Ontario only), mites as a post-harvest application on raspberries, and pear psylla in pears.

COMMERCIAL

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 75%

REGISTRATION NO.: 25135
PEST CONTROL PRODUCTS ACT



WARNING

POISON

Warning, contains the allergen sulfite

**DANGER: FATAL IF INHALED. DO NOT BREATHE DUST, VAPOUR OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES OR CLOTHING.**

**HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN.
READ THE LABEL AND BOOKLET BEFORE USING**

DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY

NET CONTENTS:

Gowan
CANADA

Canyon Group, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

NEXTER® is a registered trademark of Nissan Chemical Industries Limited used under license by Gowan Company, L.L.C.

GROUP	21	INSECTICIDE
-------	----	-------------

NEXTER® WP Miticide/Insecticide

For control of mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards (Ontario only), mites as a post-harvest application on raspberries, and pear psylla in pears

COMMERCIAL

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 75%

REGISTRATION NO.: 25135
PEST CONTROL PRODUCTS ACT



WARNING

POISON

Warning, contains the allergen sulfite

**DANGER: FATAL IF INHALED. DO NOT BREATHE DUST, VAPOUR OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES OR CLOTHING.**

**HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN.
READ THE LABEL AND BOOKLET BEFORE USING**

DO NOT SELL WATER-SOLUBLE BAGS SEPARATELY

NET CONTENTS:



Canyon Group, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

1. May be harmful if swallowed. Do not take internally.
2. May be fatal if inhaled. Do not inhale dust, vapour or spray mist. May irritate eyes. Avoid contact with eyes, skin and clothing.
3. Wash thoroughly after handling and before eating, drinking or smoking.
4. Wear long-sleeved shirt, long pants, rubber boots, socks, goggles, gloves (rubber, PVC, neoprene or nitrile), hat and a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides during mixing, loading and application. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI). Employers should make every effort to schedule pesticide applications and worker tasks in order to avoid early entry of workers into treated areas. Under exceptional circumstances, certified pesticide applicators may enter treated areas for short-term tasks not involving hand labour if at least 4 hours have passed since application and a long sleeved shirt, long pants, rubber boots, socks, goggles, gloves (rubber, PVC, neoprene or nitrile), hat, and a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 24 hour period or until the REI ends.
5. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
6. Clean spray equipment thoroughly after use.
7. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
8. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply or allow **NEXTER WP Miticide/Insecticide** to drift to crops or weeds on which bees are actively foraging. Do not allow spray to drift towards beehives.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca .

This product is only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the pesticide application is to occur.

DO NOT use in residential areas. Residential areas are defined as sites where bystanders including children may be potentially exposed during or after spraying. This includes around homes, school, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed.

Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

"Hazardous to humans and domestic animals. Keep out of reach of children and pets."

DO NOT allow entry or allow workers to enter treated areas during the restricted entry intervals REI(s) specified in the following table.

Crop	Postapplication Activity	Restricted Entry Interval
Grapes	Girdling, turning	54 Days
	Tying (full foliage), training (full foliage), hand harvest, leaf pulling by hand	30 days
	All other activities (e.g. scouting, weeding, irrigation, bird control, propagating, trellis repair, pruning, transplanting)	12 hours
All other crops	All activities	12 hours

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a major spill, leak or fire call Chemtrec® at 1-800-424-9300. For medical emergencies call 1-888-478-0798

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

1. Store in original tightly closed container. Do not expose water-soluble bags to moisture.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.

DISPOSAL

1. Follow provincial instructions for any required cleaning of the container prior to its disposal.

2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Nexter – WSB



NEXTER® WP MITICIDE/INSECTICIDE
COMMERCIAL

Pyridaben 75% Wettable Powder

REGISTRATION NO. 25135
PEST CONTROL PRODUCTS ACT

NET CONTENTS:



WARNING

POISON

Warning, contains the allergen sulfite.

KEEP OUT OF REACH OF CHILDREN.
READ THE LABEL AND BOOKLET BEFORE USING.

Product Information: 1-800-883-1844