Green Earth® BORDO COPPER SPRAY

Wettable Powder •Fungicide Spray

DOMESTIC

REG. NO. 17482 P.C.P ACT

GUARANTEE: Copper, present as basic copper sulfate 53%

WARNING



READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

200 g

Sure-Gro IP Inc. / Lic. Premier Tech Home & Garden Inc. P.O. Box 21001, Brantford, ON N3R 7W9 1-800-268-2806 Controls anthracnose, blights, leaf spots, mildew and molds on fruits, vegetables and ornamentals.

DIRECTIONS FOR USE: Green Earth® BORDO Copper Spray is a wettable powder with improved suspension in water to be used as a spray for control of certain plant diseases. Mix with water thoroughly as directed beside and keep the mixture agitated while spraying. Spray to thoroughly wet all foliage and stems. May be applied (with a Hose End Sprayer or a tank sprayer) up to 1 day before harvest. May stain paint, stone or brick. DO NOT apply this product in a way that will contact adults, children or pets, either directly or through drift. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. DO NOT apply to any body of water. Avoid application of this product when winds are gusty.

NOTE: 5 ml (1 level teaspoon) of Green Earth[®] BORDO Copper Spray weighs 3.33 g. The amount of product indicated in the opposite table is recommended to be mixed with 1 L of water and applied over a minimum area of 10 m². Adjust rates according to use area coverage.

Maximum Single Application Rate (teaspoon of product $/ 10 \text{ m}^2$) = M.S.R.

Minimum Application Interval (days) = M.A.I. (Minimum number of days between each application)

Maximum Number of Applications per Year = M.P.Y.

Use site	Diseases Controlled	Specific Directions	M.S.R	M.A.I.	M.P.Y
Apple	Anthracnose		1	10	21
Peach	Anthracnose, Coryneum Blight, Peach Leaf Curl, Shot Hole	Dormant spray, especially in late fall - never when trees are in leaf.	1/2	7	2
Potato	Early and Late Blight, Anthracnose, Leaf Mold, Septoria Leaf Spot.	Apply when plants are 12 to 18 cm high	1	5	10
Currant, gooseberry	Anthracnose, <i>Phyllosticta</i> and <i>Septoria</i> leaf spots, Powdery Mildew	Spray just before blossoms open and again when most of the petals have fallen.	1 1/3	10	4
Strawberry	Leaf spot.	Spray one month after planting	1	7	5
Bean	Anthracnose, Bacterial Leaf Spot, Downy Mildew	Apply according to Provincial Spray Calendar recommendations or on appearance of damage caused by disease	1 1/2	7	6

Beet	Cercopora Leaf Spot	Apply according to Provincial Spray calendar recommendations or on appearance of damage caused by disease	1 1/8	7	6
Carrot	Cercospora Leaf Spot	Apply according to Provincial Spray calendar recommendations or on appearance of damage caused by disease	1 1/8	7	5
Cabbage, broccoli, cauliflower	Downy Mildew	Spray on appearance of disease	1 1/8	7	5
Celery	Early and Late Blight, Anthracnose, Leaf Mold, Septoria Leaf Spot.	Apply when plants are 12 to 18 cm high	1	5	5
Cucumber	Angular Leaf Spot, Anthracnose, Downy Mildew, Scab.	Begin treatment when plants begin to vine	1	7	5
Pumpkin, squash, watermelon, melon	Angular Leaf Spot, Anthracnose, Downy Mildew, Scab.	Begin treatment when plants begin to vine	3/4	5	5
Eggplant, Pepper	Early and Late Blight, Anthracnose, Leaf Mold, Septoria Leaf Spot.	Apply when plants are 12 to 18 cm high	1 1/8	3	10
Spinach	Downy Mildew, White Rust		1/2	7	5
Tomato	Early and Late Blight, Anthracnose, Leaf Mold, Septoria Leaf Spot.	Apply when plants are 12 to 18 cm high	1 1/4	3	10
Grape	Downy Mildew, Black Rot, Dead Arm	Spray before blossoms open. Do not use on French hybrid or Vinifera varieties.	1 1/2	3	7
Ornamental Shrubs	Leaf spots, Black Spot, Powdery and Downy Mildew		1/4	7	10
Roses	Leaf spots, Black Spot, Powdery and Downy Mildew		1 1/2	7	2
٦					

Make a maximum of two applications per year. Make the first application after petal fall, and the second immediately after harvest

ENVIRONMENTAL HAZARDS: TOXIC to birds, small wild mammals, aquatic organisms and non-target terrestrial plants. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

PRECAUTIONS: Store in cool, dry ventilated place away from feeds and foods. May irritate eyes, nose, throat and skin and discolour skin. Avoid contact with these areas and avoid breathing dust or spray mist. Do not contaminate any body of water. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. DO NOT allow adults, children or pets to enter the treated area until sprays have dried.

FIRST AID: If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

DISPOSAL: Dispose of empty container in household garbage.

For additional product information or in case of emergency, spill or fire, call toll-free 1-800-268-2806. www.pthomeandgarden.com

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.